

SUTAB INSTRUCTIONS

Start Linzess	/2mcg	145mcg	290mcg oi	า		Take one
cansule 30 m	ninutes bef	ore breakf	ast on an em	nty stoma	ach will full g	lass of water
•			ast on an em	pry storric	2011 11111 1211 8	iass of water
and continue	e for 3 days	S.				
*Clear liquids	ALL DAY the	day before	your procedur	e on:		
			, can process			
NO SOLID FO	OODS THIS D	DAY				
milk), bouillon	, broth, app	le juice, whi	ubstance. Exai te grape juice, ithout fruit) ar	ANY soft d		(no creamer or d, Gatorade.
NO red or purp cream!!	ple!! NO ora	nge juice or	other pulp-cor	ntaining jui	ces, tomato ju	ice, milk or ice
On	@	2 PM	take 4 Dulcol	ax tablets.	Then drink at	least 8oz clear
liquids every h	our after tha	ıt.				
with 16oz of co	ool water (To	Fill Line). Sv	Open 1s bott wallow each ta is should take 1	blet with a	sip of water ar	nd then drink the
*Continue drir	nking clear li	quids throu	ghout the day!	*		
On	@		Open 2nd bot	tle of 12 ta	blets. Fill prov	ided container
with 16oz of co	ool water (To	Fill Line). Sv	wallow each ta	blet with a	sip of water ar	d then drink the
entire amount	of water in	the cup. (Th	is should take 1	15-20 minu	tes to complet	e).
NOTHING TO [ORINK AFTER	R		(4 h	rs before arriv	al time)
Report to:	9 Physicians	Drive. Jacks	on TN at	on		•

- Bring a DRIVER. Also bring your insurance card, photo ID, and a list of all of your medications.
- Notify the doctor if on any blood thinners.
- Take any blood pressure, heart, or seizure medications the morning of your procedure. DO NOT take any diabetic medications.
- If you have any questions call 731-661-0086